Contribution ID: 414

Type: Oral talk (15 min + 5 min questions)

PHOTONUCLEAR METHOD FOR THE PRODUCTION OF MEDICAL RADIOISOTOPE 72AS

Friday, 15 July 2022 15:20 (10 minutes)

The method of induced activity was used to study photonuclear reactions on a natural mixture of selenium isotopes. The experiment was performed on a bremsstrahlung of an RM-55 electron accelerator at an electron energy of 55 MeV. The study examined the possibility of producing 72As isotope in photonuclear reactions on a natural mixture of selenium isotopes. Experimental data on the cross-sections of photoproton reactions on Se isotopes are not available in the literature. The yields of the formation of 73,74,75,76,81,81mSe isotopes as a result of natSe(γ , in) reactions, the target nuclide 72As and the side nuclides 71,74,76,77,78,79As as a result of natSe(γ , in1p) reactions were measured. The experimentally obtained yields of photonuclear reactions are compared with the yields calculated using theoretical cross-sections of photonuclear reactions from and the TALYS program.

The speaker is a student or young scientist

Yes

Section

1. Nuclear technology and methods in medicine, radioecology

Primary authors: Mr ALIEV, R; BELYSHEV, Sergey (Faculty of Physics, Lomonosov Moscow State University, Russia. Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University, Russia.); Mrs FURSOVA, N (Lomonosov Moscow State University, Department of Physics); Dr KHANKIN, Vadim (Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University); KUZNETSOV, Alexander (Lomonosov Moscow State university, Faculty of Physics; Skobeltsyn Institute of Nuclear Physics, Lomonosov Moscow State University); RASULOVA, Fazilat (Institute of Nuclear Physics of AS RUz)

Presenter: RASULOVA, Fazilat (Institute of Nuclear Physics of AS RUz)

Session Classification: Nuclear technology and methods in medicine, radioecology.