

ATOMIC K-SHELL DOUBLE HOLE CREATION DUE TO ELECTRON CAPTURE AND PHOTOIONIZATION

Wednesday, 13 July 2022 16:00 (20 minutes)

Double core hole (DCH) states could be created in two different mechanisms. The first is core subshell electron capture by the nucleus and subsequent shake-process of an electron from the same subshell either to the high unoccupied state (shake-up) or to the continuum (shake-off) mostly due to instant changing of the nucleus charge by unity. Another way is photoionization, when one electron is "knocked out" from the subshell and the residual one undergoes shake-process resulting from changing in screened potential. Physics of the DCH currently attracts interest stimulated by new possibilities of their creation by X-ray free electron lasers and advanced synchrotron sources. Furthermore, DCHs might become a new tool for chemical analysis [1] and plasma diagnostics [2].

In this theoretical contribution we compare the two mechanisms of producing the DCH: K-electron capture and K-shell photoionization. General theoretical approaches to both problems are known, but we are not aware of such a comparison based on up-to-date models for many-electron atoms. We focus on DCH states in K-shell of ${}^7\text{Be}$ and ${}^{37}\text{Ar}$ (isotopes with natural electron capture radioactivity). The goal is to determine double K-vacancy production probability in both described channels. Also, we analyze shake-off electron spectra and compare the results with different theoretical approaches and experiment.

Our model is based on sudden approximation with the use of fully non-orthogonal sets of electron orbitals in initial and final states, accounting for j-splitting of the subshells. Expansions for transition matrix elements are obtained with ZAP_NO package [3]. Radial wave functions are constructed within the multiconfigurational Hartree-Fock method [4]. Photoionization calculations are performed with the use of B-spline R-matrix (BSR) software complex [5].

This research was funded by the Russian Ministry of Science and Education grant No. 075-15-2021-1353. The work of M.D.K. is supported by the Ministry of Science and Higher Education of the Russian Federation (project No. 0818-2020-0005) using resources of the Shared Services "Data Center of the Far-Eastern Branch of the Russian Academy of Sciences".

1. M. Nakano et al., Phys. Rev. Lett. 110, 163001 (2013).
2. A.Ya. Faenov et al., Laser and Particle Beams 33, 27 (2015).
3. O. Zatsarinny, Comput. Phys. Comm. 98, 235 (1996).
4. C.F. Fischer, T. Brage and P. Jonsson, Computational Atomic Structure: An MCHF Approach (Bristol: IOP Publishing, 1997).
5. O. Zatsarinny, Comput. Phys. Comm. 174, 273 (2006).

The speaker is a student or young scientist

Yes

Section

1. Synchrotron and neutron radiation sources and their use in scientific and applied fields

Primary authors: Mr KISELEV, Maksim (Lomonosov Moscow State University); Dr GRYZLOVA, Elena (Skobel'syn Institute of Nuclear Physics, Lomonosov Moscow State University); Dr GRUM-GRZHIMAILO, Alexei (Skobel'syn Institute of Nuclear Physics, Lomonosov Moscow State University)

Presenter: Mr KISELEV, Maksim (Lomonosov Moscow State University)

Session Classification: Synchrotron and neutron radiation sources and their use in scientific and applied fields