Contribution ID: 262

Type: Oral talk (15 min + 5 min questions)

Microscopic description of isoscalar giant monopole resonance: the case of 48Ca

Thursday, 14 July 2022 15:20 (20 minutes)

A comprehensive analysis of the isoscalar giant monopole resonance (ISGMR) has long been a subject of extensive theoretical and experimental research [1,2]. The ISGMR properties are presently an important problem not only from the nuclear structure point of view [2,3] but also because of the special role they play in many astrophysical processes such as prompt supernova explosions [4] and the interiors of neutron stars [5]. One of the successful tools for describing the ISGMR is the quasiparticle random phase approximation (QRPA) with the self-consistent mean-field derived from Skyrme energy density functionals (EDF) [2,3]. The study of the monopole strength distribution in the region of giant resonance involves taking into account a coupling between the simple particle-hole excitations and more complicated (two- and three-phonons) configurations [3,6]. The main difficulty is that the complexity of calculations beyond standard QRPA increases rapidly with the size of the configuration space, and one has to work within limited spaces. Using a finite rank separable approximation for the residual particle-hole interaction derived from the Skyrme EDF one can overcome this numerical problem [7,8].

In the present report, we study the effects of the coupling between one-, two- and three-phonon terms in the wave functions on the monopole strength distribution in the double magic nucleus 48Ca. Using the same set of parameters, we describe available experimental data [9,10]. The effects of the phonon-phonon coupling leads to a redistribution of the main monopole strength to lower energy states and also to higher energy tail [11].

This work was supported by the Russian Science Foundation (Grant No. RSF-21-12-00061).

- 1. J.P. Blaizot, Phys. Rev. 64, 171 (1980).
- 2. U. Garg, and G. Colò, Prog. Part. Nucl. Phys. 101, 55 (2018).
- 3. N.N. Arsenyev, and A.P. Severyukhin, Universe. 7, 145 (2021).
- 4. H.A. Bethe, Rev. Mod. Phys. 62, 801 (1990).
- 5. N.K. Glendenning, Phys. Rev. Lett. 57, 1120 (1986).
- 6. V.G. Soloviev, Theory of Atomic Nuclei: Quasiparticles and Phonons. 1992. Bris-tol/Philadelphia.
- 7. N.V. Giai, Ch. Stoyanov, and V.V. Voronov, Phys. Rev. C. 57, 1204 (1998).
- 8. A.P. Severyukhin, V.V. Voronov, and N.V. Giai, Eur. Phys. Jour. A. 22, 397 (2004).
- 9. K. Howard et al., Phys. Lett. B. 801, 135185 (2020).
- 10. S.D. Olorunfunmi et al., arXiv:2202.00722v1 [nucl-ex] 1 Feb 2022.
- 11. N.N. Arsenyev, and A.P. Severyukhin, in preparation.

The speaker is a student or young scientist

No

Section

1. Nuclear structure: theory and experiment

Primary authors: Dr ARSENYEV, Nikolay (Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research); SEVERYUKHIN, Alexey (BLTP, JINR)

Presenter: Dr ARSENYEV, Nikolay (Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research)

Session Classification: Nuclear structure: theory and experiment