

Pion femtoscopy in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV in the STAR experiment

Tuesday, 12 July 2022 10:20 (20 minutes)

Pion femtoscopy in Au+Au collisions at $\sqrt{s_{NN}} = 3$ GeV in the STAR experiment

A. Kraeva (for the STAR Collaboration)

E-mail: annakraeva555@gmail.com

National Research Nuclear University MEPhI, Kashirskoe highway 31, Moscow, 115409, Russia

There is a method that allows directly measuring the spatio-temporal extent of the region where hadrons are emitted and the parameters of the nuclear-nuclear interaction, called femtoscopic correlation [1]. In heavy-ion collisions, femtoscopy is an important tool for studying the geometric and dynamic characteristics of the emission region.

Two-particle momentum correlations of identical particles in nuclear-nuclear collisions make it possible to extract femtoscopic parameters (radii of emission region, R , and correlation strength, λ) [2]. Reaction dynamic is reflected in the femtoscopic radii dependence on pair transverse momentum, k_T .

This work is devoted to the study of two-particle momentum correlations of identical pions produced in collisions of gold nuclei in the STAR experiment at the RHIC at $\sqrt{s_{NN}} = 3$ GeV. The extracted three-dimensional femtoscopic radii (R_{out} , R_{side} , R_{long}) are measured as a function of collision centrality and transverse momentum of the pairs.

References:

[1] Podgoretsky M.I. Interference correlations of identical pions, Sov. J. Nucl. Phys. – 1989. – V.20. – P.3.

[2] Lisa M.A. et al. Femtoscopy in relativistic heavy ion collisions: two decades of progress, Annu. Rev. Nucl. Part. Sci. – 2005. – V.55. – P.357.

The speaker is a student or young scientist

Yes

Section

1. Intermediate and high energies, heavy ion collisions

Primary author: KRAEVA, Anna

Presenter: KRAEVA, Anna

Session Classification: Intermediate and high energies, heavy ion collisions