Contribution ID: 113

Type: Oral talk (15 min + 5 min questions)

First measurement of the forward rapidity gap distribution in proton-lead collisions at LHC energy $\sqrt{s_{\mathrm{NN}}} = 8.16~\mathrm{TeV}$ with the CMS experiment

Tuesday, 12 July 2022 15:00 (20 minutes)

For the first time at LHC energies, we present the forward rapidity gap spectra from proton-lead (pPb) collisions for both pomeron-lead and pomeron-proton topologies. The analysis is performed over 10.4 units of pseudorapidity. The center-of-mass energy is 8.16 TeV per nucleon-nucleon pair, i.e., almost 300 times higher than previous measurements of diffractive production in proton-nucleus collisions. For the pomeron-lead topology, the EPOS - LHC predictions are a factor of two below the unfolded data, but the model does give a reasonable description of the spectrum's shape. For the pomeron-proton topology, the EPOS-LHC, QGSJET II, and HIJING generator predictions are all at least a factor of five below the data. This effect can be explained by a significant contribution of ultra-peripheral photoproduction events mimicking the signature of diffractive processes. The obtained data may be of significant help in understanding the high energy limit of QCD and modeling cosmic ray air showers.

The speaker is a student or young scientist

No

Section

1. Intermediate and high energies, heavy ion collisions

Primary author: SOSNOV, Dmitry (NRC KI – PNPI)

Presenter: SOSNOV, Dmitry (NRC KI – PNPI)

Session Classification: Intermediate and high energies, heavy ion collisions