

ON THE WIDTH OF γ -LINE AND THE PHOTON STRUCTURE

Wednesday, 13 July 2022 11:10 (20 minutes)

The reason to turn once again to the question of the natural width Γ_γ of the γ -radiation line of nuclei is provided by the discovery of an increase in the half-life $T_{1/2}$ for nuclear isomers in a metal matrix (see ref. [1] and references therein). For nuclei in such a matrix, a decrease in the width Γ_γ can be expected, since, according to generally accepted concepts, $\Gamma_\gamma \propto 1/T_{1/2}$ for the emission of photons in a nuclear transition from an excited state E of a half-life $T_{1/2}$ to the ground state.

However, even in early Mössbauer experiments with the ^{57}mFe isomer, a decrease in Γ_γ was observed with an increase in the age of the E level (see, e.g., ref. [2]), which could be interpreted as a result of a decrease in the level width with its age. But such an interpretation is not allowed by the experiment [3] with the $^{181\text{m}}\text{Ta}$ isomer ($T_{1/2} \approx 6 \mu\text{s}$), in which broadening of the 6.2 keV γ -line was observed due only to the shading of the absorber from the emitter by a mechanical chopper, which opened their mutual visibility for a time of 1 μs without referencing by the time the isomer was formed. Hence it follows that the width Γ_γ is determined not by the value of $T_{1/2}$, but only by the time T_γ , which in the Mössbauer experiments the absorber nucleus sees the emitter before the emission of an energy quantum. Of course, if there are no restrictions on the measurement time of the width Γ_γ , then the average value T_γ is proportional to $T_{1/2}$.

Then, taking into account that the energy of the γ -transition is emitted in less than 1 ns – this can be seen, for example, from the duration of the γ -signal in the scintillator, we can assume the following photon structure. Immediately after the formation of the excited state E , the nucleus begins to emit an electromagnetic wave of frequency ω that does not carry energy – abbreviated as a 0-wave. The duration of this 0-wave determines the width Γ_γ . The energy quantum $\hbar\omega$ is emitted at the end of the 0-wave. The energy $\hbar\omega$ may not be emitted at all if the state E decays via another channel, and then the 0-wave will exist on its own, without an energy quantum. A possible source of the 0-wave is the virtual transitions from the E^* level to the ground state and back before the emission of an energy quantum.

The 0-wave with a quantum $\hbar\omega$ "on its tail" resembles a pilot wave introduced by De Broglie to explain the wave-particle duality of electrons. It is interesting to study effect of 0-waves on absorber nuclei, for example, to search for the modulation of the Γ_γ value via an additional resonance irradiation of the absorber in Mössbauer experiments.

1. V.V. Koltsov, Bull. Russ. Acad. Sci.: Phys. 83, 1144 (2019).
2. W. Triftshausen, P.P. Craig, Phys. Rev. Lett. 16, 1161 (1966).
3. V.K. Voitovetsky, I.L. Korsunsky, Yu.F. Pazhin et al., Yad. Fiz. 38, 662 (1983).

The speaker is a student or young scientist

No

Section

1. Experimental and theoretical studies of nuclear reactions

Primary author: KOLTSOV, Vladimir (Khlopin Radium Institute, Saint-Petersburg, Russian Federation)

Presenter: KOLTSOV, Vladimir (Khlopin Radium Institute, Saint-Petersburg, Russian Federation)

Session Classification: Poster session