Contribution ID: 494

Type: Oral talk (15 min + 5 min questions)

ANALYSIS OF THE COMPOSITION OF ISOTOPES IN A SAMPLE OF HIGH-TEMPERATURE SUPERCONDUCTORS (HTS) AFTER IRRADIATION IN THE RADIATION FIELD OF A LEAD CONVERTER WITH A PROTON BEAM (EP = 660 MeV)

Friday, 15 July 2022 18:10 (20 minutes)

Combined talk

ANALYSIS OF THE COMPOSITION OF ISOTOPES IN A SAMPLE OF HIGH-TEMPERATURE SUPERCON-DUCTORS (HTS) AFTER IRRADIATION IN THE RADIATION FIELD OF A LEAD CONVERTER

WITH A PROTON BEAM (EP = 660 MeV)

M.S. Novikov, S.I. Tyutyunnikov, G.V. Trubnikov, V.I. Stegailov,

V.P.Volnykh, T.N. Tran.

JINR, Dubna, Russia, E-mail: stegajlov2013@yandex.ru, tntoan@iop.vast.vn

This report considers one of the stages of work on the development of a series of superconducting magnets operating at temperatures up to 65 K for the possible creation on the basis of a series of accelerators: proton cyclotron (Phasotron), medical cyclotron (SC230 with HTS winding, 4kA, 2-T) and etc.

1. Novikov M., Khodzhibagiyan H., Tyutyunnikov S., Stegailov V., Smirnov G., Petrova M., Kondratiev B. Book of abstracts and articles of Int. Conf. "Modern Problems of Nuclear Energetics and Nuclear Technologies". P. 111. Tashkent, November (2021).

ION SOURCE FOR RESEARCH IN NUCLEAR PHYSICS

V.P. Ovsyannikov1, A.V. Nefedov2, A.A. Levin3, V.I. Stegailov1, S.I. Tyutyunnikov1, Y.A.Vaganov1

1Joint Institute for Nuclear Research, 141980 Dubna, Russia; 2Petersburg Nuclear Physics Institute, 188300 Gatchina, St. Petersburg, Russia; 3Ioffe Institute, 194021 St. Petersburg,

E-mail: stegajlov2013@yandex.ru, vaganov ya@jinr.ru

An ionizer of the MaMFIS type (Main Magnetic Focus Ion Source) [1] produces ions with a high charge in local ion traps of a pulsating electron beam, where the density can still reach extremely high values (~10 kA/cm2). X-ray radiation from Bi60+ ions was detected. The peculiarity of the ionizer - a high ion charge and a relatively low number of ions, makes it attractive for experiments with an ion Penning trap (MATS and LaSpec: High-precision experiments using ion traps and lasers at FAIR).

1. V.P. Ovsyannikov, A.V. Nefiodov, A.Yu. Boytsov, A.Ya. Ramsdorf, V. I. Stegailov, S.I. Tyutyunnikov, A.A. Levin // Main magnet, NIM, 502 (2021) 23-28.

152Eu NUCLEUS IN THE FIELD OF LASER RADIATION 1064nm E.V. Barminal, A.V. Simakin1, S.I. Tyutyunnikov2, V.I. Stegailov2, G.A. Shafeev1,3, I.A. Shcherbakov1

1Prokhorov General Physics Institute of the Russian Academy of Sciences, Moscow, Russia; 2Joint Institute for Nuclear Research, Dubna, Russia; 3National Research Nuclear University "MEPHI", Moscow, Russia E-mail: stegajlov2013@yandex.ru

The experiments were carried out within the framework of the project «Energy and Transmutation of Radioactive Waste». The decay of nuclei under the action of laser radiation with a wavelength of 1064 nm on their aqueous solutions was studied. The well-studied 152Eu nucleus is of particular interest. 1. E.V. Barmina, S.I.Tyutyunnikov et al. Quantum Electronics (2019), 49 (8):784.

The speaker is a student or young scientist

No

Section

1. Experimental and theoretical studies of nuclear reactions

Primary author: Mr STEGAYLOV, Vladimir (JINR)

Presenter: Mr STEGAYLOV, Vladimir (JINR)

Session Classification: Experimental and theoretical studies of nuclear reactions